

## Advil Cold & Sinus Caplets & Liqui-Gels®



### Active Ingredients

- Ibuprofen 200mg
- Pseudoephedrine hydrochloride 30mg

### Uses

For effective relief of the symptoms of colds and sinusitis including:

- nasal congestion
- body aches and pain
- sinus pain
- sore throat
- headache
- fever

### Directions

Adults and children over 12: Take 1 or 2 caplets or liqui-gels every 4 hours as needed. Do not exceed 6 caplets or liqui-gels in 24 hours, unless directed by a physician.

### • Caution

Keep out of reach of children. This package contains enough medicine to seriously harm a child. Do not give to children under 12 unless directed by a physician.

### Warning

Do not take Advil Cold & Sinus if taking acetylsalicylic acid (ASA) or other products containing ibuprofen, or if allergic to ASA, salicylates or anti-inflammatory drugs, or any of its ingredients. Consult your physician before taking Advil Cold & Sinus if you have kidney or liver disease, diabetes, heart or thyroid disease, high blood pressure, glaucoma, peptic ulcers, difficulty in urination due to enlarged prostate, or if you are pregnant or nursing a baby. Individuals taking or who have recently taken prescription medication for high blood pressure or depression, including monoamine oxidase inhibitors (MAOIs), should use only as directed by a physician. Consult your physician if the symptoms or fever persists for more than 5 days.

If abdominal pain, heartburn, nausea or vomiting, bloating, diarrhea or constipation, ringing or buzzing in the ears, nervousness, sleeplessness, dizziness, any change in vision, fluid retention, itching, skin rashes or any other side effect or unexplained symptom develops while taking Advil Cold & Sinus, discontinue use immediately and contact a physician. In case of overdose, call Poison Control Centre or a doctor at once even if there are no symptoms.


### Caplets Non-medicinal Ingredients

acetylated monoglycerides, ammonium hydroxide, carnauba wax, corn starch, croscarmellose sodium, ethoxyethanol, iron oxides, lecithin, microcrystalline cellulose, parabens, pharmaceutical glaze, pharmaceutical shellac, povidone, pregelatinized starch, propylene glycol, silicon dioxide, simethicone, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, titanium dioxide.

### Liqui-Gels Non-medicinal Ingredients

D&C yellow no. 10, FD&C red no. 40, fractionated coconut oil, gelatin, iron oxide, polyethylene glycol, polyvinyl acetate phthalate, potassium hydroxide, propylene glycol, purified water, sorbitan, sorbitol.

## Advil Cold & Sinus PLUS



### Active Ingredients

- Ibuprofen 200mg
- Pseudoephedrine hydrochloride 30mg
- Chlorpheniramine maleate 2mg

### Uses

For effective relief of the symptoms of colds and sinusitis including:

- nasal congestion
- fever
- body aches and pains
- sinus pain
- runny nose
- sore throat
- headache
- sneezing
- itching of the throat

### Directions

Adults and children over 12: Take 1 or 2 Advil Cold & Sinus PLUS caplets every 4 hours as needed. Do not exceed 6 Advil Cold & Sinus PLUS caplets in 24 hours, unless directed by a physician.

### • Caution

Keep out of reach of children. This package contains enough medicine to seriously harm a child. Do not give to children under 12 unless directed by a physician.

### Warning

Do not take Advil Cold & Sinus PLUS if taking acetylsalicylic acid (ASA) or other products containing ibuprofen, or if allergic to ASA, salicylates or anti-inflammatory drugs, or any of its ingredients. Consult your physician before taking Advil Cold & Sinus PLUS if you have kidney or liver disease, diabetes, heart or thyroid disease, chronic lung disease, high blood pressure, glaucoma, peptic ulcers, difficulty in urination due to enlarged prostate, or if you are pregnant or nursing a baby. Individuals taking or who have recently taken prescription medication for high blood pressure or depression, including monoamine oxidase inhibitors (MAOIs), other antihistamines, tranquilizers or sedating drugs should use only as directed by a physician. Consult your physician if the pain or fever persists for more than 5 days. Product may cause drowsiness. Avoid alcoholic beverages, driving a motor vehicle or operating machinery. Do not exceed the recommended dose unless advised by a physician.

If abdominal pain, heartburn, nausea or vomiting, bloating, diarrhea or constipation, ringing or buzzing in the ears, nervousness, sleeplessness, dizziness, any change in vision, fluid retention, itching, skin rashes or any other side effect or unexplained symptom develops while taking Advil Cold & Sinus PLUS, discontinue use immediately and contact a physician. In case of overdose, call a Poison Control Centre or a doctor at once even if there are no symptoms.

### Non-medicinal Ingredients

carnauba wax, corn starch, croscarmellose sodium, FD&C red no. 40, FD&C yellow no. 6, glyceryl behenate, hypromellose, iron oxide, microcrystalline cellulose, polydextrose, polyethylene glycol, pre-gelatinized starch, propylene glycol, silicon dioxide, titanium dioxide.

## Advil Cold & Sinus Day & Night Convenience Pack



### Active Ingredients

- Daytime Caplets:*
- Ibuprofen 200 mg
  - Pseudoephedrine hydrochloride 30 mg
- Nighttime Caplets:*
- Ibuprofen 200 mg
  - Pseudoephedrine hydrochloride 30 mg
  - Chlorpheniramine Maleate 2 mg

### Uses

#### Advil Cold & Sinus Daytime

For effective relief of the symptoms of colds and sinusitis including:

- nasal congestion
- fever
- sinus pain
- sore throat
- headache
- itchy, watery eyes

#### Advil Cold & Sinus Nighttime: PLUS relief of:

- runny nose
- sneezing

### Directions

Adults and children over 12: Take 1 or 2 Advil Cold & Sinus Daytime or Nighttime caplets every 4 hours as needed. Do not exceed a combined total of any 6 caplets of Advil Cold & Sinus Daytime or Nighttime caplets in 24 hours, unless directed by a physician.

### • Caution

Keep out of reach of children. This package contains enough medicine to seriously harm a child. Do not give to children under 12 unless directed by a physician.

### Warning

Do not use if taking acetylsalicylic acid (ASA) or other products containing ibuprofen, or if allergic to ASA, salicylates or anti-inflammatory drugs, or any of its ingredients. Consult your physician before taking Advil Cold & Sinus Daytime or Nighttime if you have kidney or liver disease, diabetes, heart or thyroid disease, chronic lung disease, high blood pressure, glaucoma, peptic ulcers, difficulty in urination due to enlarged prostate, or if you are pregnant or nursing a baby. Individuals taking or who have recently taken prescription medication for high blood pressure or depression, including monoamine oxidase inhibitors (MAOIs) should use only as directed by a physician. Advil Cold & Sinus Nighttime caplets may cause drowsiness and if you are taking other antihistamines, tranquilizers or sedating drugs, consult your doctor. Avoid alcoholic beverages, driving a motor vehicle or operating machinery. In case of overdose, call a poison control centre or a doctor at once even if there are no symptoms. Consult your physician if the symptoms or fever persists for more than 5 days.

### Daytime Non-medicinal Ingredients

acetylated monoglycerides, ammonium hydroxide, carnauba wax, corn starch, croscarmellose sodium, ethoxyethanol, iron oxides, lecithin, microcrystalline cellulose, parabens, pharmaceutical glaze, pharmaceutical shellac, povidone, pregelatinized starch, propylene glycol, silicon dioxide, simethicone, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, titanium dioxide.

### Nighttime Non-medicinal Ingredients

carnauba wax, corn starch, croscarmellose sodium, FD&C red no. 40, FD&C yellow no. 6, glyceryl behenate, hypromellose, iron oxide, microcrystalline cellulose, polydextrose, polyethylene glycol, pre-gelatinized starch, propylene glycol, silicon dioxide, titanium dioxide.